

Comment on “Lyapunov Exponent of a Many Body System and Its Transport Coefficients”

In a recent Letter, Barnett, Tajima, Nishihara, Ueshima, and Furukawa [1] obtained a theoretical expression for the maximum Lyapunov exponent λ_1 of a dilute gas. They conclude that λ_1 is proportional to the cube root of the self-diffusion coefficient D , independent of the range of the interaction potential. They validate their conjecture with numerical data for a dense one-component plasma, a system with long-range forces. We claim that their result is highly nongeneric. We show in the following that it does not apply to a gas of hard spheres, neither in the dilute nor in the dense phase.

Systems of hard spheres have properties similar to real fluids and solids and provide a reference for successful perturbation theories [2]. Simulations with this model were able to uncover fundamental aspects of collective particle dynamics such as recollisions and the “cage” effect [2]. Hard-sphere systems are also paradigms for the chaotic and ergodic properties of many body systems with short-range interactions, and were shown to have a positive Kolmogorov-Sinai entropy [3,4].

For dilute gases, Krylov [5] provided an analytical estimate for the maximum Lyapunov exponent,

$$\lambda_1 = -(32\pi K/3mN)^{1/2} \sigma^2 n \log(\pi \sigma^3 n / \sqrt{2}), \quad (1)$$

where K is the kinetic energy, N is the number of particles, m is the particle mass, n is the number density, and σ is the hard-sphere diameter. This expression has been verified numerically (apart from a factor ~ 2.8 [4]), and has been extended to larger densities [6].

The diffusion coefficient for dilute hard-sphere gases is well approximated by the Enskog expression [7],

$$D_E = (3\pi K/32mN)^{1/2} \frac{1}{n\pi\sigma^2} \left[1 + \frac{5n\pi\sigma^3}{12} \right]^{-1}. \quad (2)$$

A comparison of Eqs. (1) and (2) reveals that, in the dilute gas limit, the proposed relation $\lambda_1 \propto D^{1/3}$ of Barnett *et al.* cannot be satisfied. Moreover, we combine in Fig. 1 recent simulation results for D and λ_1 , which were obtained for a system of 500 hard spheres over the full range of fluid densities ($0.0001 < n\sigma^3 < 0.89$). Reduced units are used for which σ , m , and the kinetic energy per particle K/N are all unity. One observes that these data are not consistent with the proposed $D^{1/3}$ dependence (solid line), neither for low densities nor for large.

We conclude that the conjecture by Barnett *et al.* does not apply to many body systems with short-range interactions. But even its applicability for long-range interactions is doubtful. A one-dimensional gravitational system with finite N exhibits a positive λ_1 [8], whereas this clustering and confining system does not show diffusion. We also note that, while the theoretical expression (26) in Ref. [1] has been obtained for a dilute gas, the




FIG. 1. Simulation results $\lambda_1 = \lambda_1(D)$ (crosses) for a gas of hard spheres. The solid line refers to the expression $\lambda_1 \propto D^{1/3}$ suggested in Ref. [1].

data in Fig. 1 of Ref. [1] are for a dense plasma with a Coulomb coupling constant Γ ranging from 1 to 150. As reported by the same authors [9], for $\Gamma > 1$ the plasma behaves as a liquid and not as a gas. The dilute gas limit is recovered only for $\Gamma \ll 1$ [9].

We thank M. Antoni, U. Balucani, A. Rapisarda, and S. Ruffo for useful discussions.

A. Torcini,¹ Ch. Dellago,² and H. A. Posch³

¹Dipartimento di Energetica
INFN, Unità di Firenze
I-50139 Firenze, Italy

²Department of Chemistry
University of California
Berkeley, California 94720

³Institut für Experimentalphysik
Universität Wien
Boltzmannngasse 5
A-1090 Wien, Austria

Received 25 March 1998

PACS numbers: 05.45.-a, 05.60.-k

- [1] D. M. Barnett, T. Tajima, K. Nishihara, Y. Ueshima, and H. Furukawa, *Phys. Rev. Lett.* **76**, 1812 (1996).
- [2] J. P. Boon and S. Yip, *Molecular Hydrodynamics* (McGraw-Hill, New York, 1980).
- [3] H. van Beijeren, J. R. Dorfman, H. A. Posch, and Ch. Dellago, *Phys. Rev. E* **56**, 5272 (1997).
- [4] Ch. Dellago and H. A. Posch, *Physica (Amsterdam)* **240A**, 68 (1997).
- [5] N. S. Krylov, *Works on the Foundations of Statistical Mechanics* (Princeton University Press, Princeton, NJ, 1979).
- [6] R. van Zon, H. van Beijeren, and Ch. Dellago, *Phys. Rev. Lett.* **80**, 2035 (1998).
- [7] S. Chapman and T. G. Cowling, *The Mathematical Theory of Non-Uniform Gases* (Cambridge University Press, Cambridge, England, 1970).
- [8] Lj. Milanović, H. A. Posch, and W. Thirring, *Phys. Rev. E* **57**, 2763 (1998).
- [9] Y. Ueshima *et al.*, *Phys. Rev. E* **55**, 3439 (1997).